



Armenian virtual observatory simple image access service



A.V. Knyazyan^{a,*}, H.V. Astsatryan^a, A.M. Mickaelian^b

^a Institute for Informatics and Automation Problems, National Academy of Sciences of the Republic of Armenia, 1, P. Sevak str., Yerevan 0014, Armenia

^b V. Ambartsumian Byurakan Astrophysical Observatory, National Academy of Sciences of the Republic of Armenia, Byurakan 0213, Aragatzotn province, Armenia

ARTICLE INFO

Article history:

Received 7 October 2016

Accepted 3 March 2017

Available online 8 March 2017

Keywords:

ArVO

IVOA

Virtual observatory

SIA

Data sharing

ABSTRACT

The aim of the article is to introduce the data sharing service of the Armenian Virtual Observatory (ArVO) based on the Simple Image Access (SIA) Protocol of the International Virtual Observatory Alliance (IVOA).

© 2017 Elsevier B.V. All rights reserved.

Contents

1. Introduction.....	23
2. Data access standards and protocols.....	23
3. ArVO SIA service.....	24
4. Scientific applications of ArVO SIA service.....	25
5. Conclusion.....	25
References.....	26

1. Introduction

Data volumes generated in astronomy are very large and observations taken at a particular time are by definition unrepeatable and irreplaceable. The overall quality of telescopes and detectors is getting better year-by-year. It is expected that some of the surveys, which will be initiated over the next few years, will generate terabytes of data per day (Gudivada et al., 2015; Mickaelian et al., 2016b).

During recent years Byurakan Astrophysical Observatory (Murdin, 2000) astronomers have done a huge work to digitize the plates of past years. This First Byurakan Survey (FBS) data is the core of the ArVO database (Mickaelian et al., 2007). It includes Digitized First Byurakan Survey, the largest and the first systematic objective prism survey of the extragalactic sky. At the same time Byurakan Astrophysical Observatory telescopes continuously generate astronomical data. At the observatory five observational instruments are installed; the larger ones being 2.6 m Cassegrain

telescope and 1 m Schmidt telescope. Storing, archiving, reducing, retrieving, analyzing, sharing and scientifically using these vast data are a formidable task. Virtual Observatories (VOs) seek to facilitate the mentioned task by using international standards and protocols.

2. Data access standards and protocols

A suite of IVOA (Quinn et al., 2004) protocols specifies basic access methods for different types of data resource, such as Simple Image Access (SIA) (Tody et al., 2009) for image archives, Simple Cone Search (Plante et al., 2008) for source catalogues, Simple Spectral Access (SSA) (Tody and Dolensky, 2004) for spectra, and several others. Only a few parameters can be involved in the data access requests (typically just RA, Dec, and search radius), and implementing these services is relatively easy. As a result all these protocols are called simple. IVOA provides Table Access Protocol (TAP) (Dowler, 2016), for more flexible access.

The Virtual Observatory (VO) is based around the two key concepts of Resources and Services (Brunner et al., 2001). “Resource” covers a wide variety of things, such as a database, a web page, a

* Corresponding author.

E-mail addresses: aram_knyazyan@ipia.sci.am (A.V. Knyazyan), hbrach@sci.am (H.V. Astsatryan), aregmick@yahoo.com (A.M. Mickaelian).




Fig. 1. Request/response process to ArVO SIA API.

storage element, and so on. All above mentioned things basically are uniquely addressable through the Internet. For example, a “data service” is a communication system that allows sending queries to a database resource and getting back data in return.

The service architecture has to follow some standard protocol, like the data access protocols described above. It also needs to publish service metadata that describes how it works, what it is and what it provides. Based on IVOA SIA protocol, a simple image search service for ArVO has been implemented. Its functionality is presented in detail in the next section.

3. ArVO SIA service

SIA protocol has 2 versions—SIA1.0 (Tody et al., 2009) and SIA2.0 (Dowler, 2015). In the first phase the implementation of ArVO SIA service was planned based on the SIA1.0 protocol. The extension of the service using the SIA2.0 is under construction. SIA provides capabilities for the discovery, description, access, and retrieval of multi-dimensional image datasets. It works using standard http get, and allows passing several standard query parameters. The output returned by an Image Query is a VOTable, an XML table format, returned with a MIME-type of text/xml. The purpose of the image query web method is to allow users to search for image data for a given region on the sky. Additional parameters may optionally be used to further refine the query. The size, scale, projection, and so forth of the ideal output image can be specified, and the service will return references to images, which most closely match what is requested. The request and response process is presented in Fig. 1.

A base URL for SIA request must be defined, to which web query parameters can be passed. The POS and the SIZE are mandatory parameters to be passed in get query. The POS is a position of the region of interest, expressed as the right ascension (RA) and declination (DEC) of the field center, in decimal degrees using the ICRS coordinate system. A comma should delimit the two values (POS = 12.713, −21.1).

The SIZE parameter presents the angular size of the region given in decimal degrees (SIZE = 1). Thus, having a base URL and passing POS and SIZE parameters will allow to construct a minimal and simple image access request.

For more complex requests it needs to use optional parameters, such as:

- NAXIS—the size of the output image in pixels. ArVO images size is predetermined and for given region of search can exist many images with different sizes. This parameter is used for returning the image whose size matches with provided NAXIS. The re-sampling mechanism of the output image size is under development.
- EQUINOX—epoch of the mean equator. EQUINOX is used for correcting the search coordinates, as ArVO data contains both B1950 and J2000 images.
- VERB—the desired level of information to be returned in the output table. Two levels of verbosity are implemented, 0 for minimal information for returned images, and 1 for regular number of columns with all available information.
- FORMAT—the desired format or formats of the images referenced by the output table, and so on. More information about optional parameters is available in SIA documentation (Tody et al., 2009).

The list of these parameters is regularly updated, and new functionalities are included in provided service. The returned table lists all the images available to the client that match the query constraints. It must contain a RESOURCE element, identified with the tag type = “results”, containing a single TABLE element which contains the results of the query. The RESOURCE element contains an INFO with name = “QUERY_STATUS”. Its value in case of success is OK and ERROR in case of any issue. It also includes all astronomical metadata of found images, their physical address URLs and so on. The result contains one record per image. Both calibrated and raw data can be returned. If calibrated data is not available, raw data will be returned. User can provide NAXIS parameter to get directly the raw data.

A simple web query example to ArVO SIA is <http://arvo.sci.am/api/search.php?POS=322.3,3&SIZE=1>. The base URL is <http://arvo.sci.am/api/search.php>, POS and SIZE query parameters are passed. The result of mentioned query is presented in Fig. 2.

SIA and other IVOA services of different VOs can be added in astronomical registries which allow discovering all published VO resources around the world. Thus, different astronomical resources become available in one place, and its data can be accessed by the same rules and request parameters. Each VO SIA service has a unique base URL, and a request with same parameters can be sent in parallel to all registered services. Hence, an astronomer will be able to get an entire part of universe observation results of different observatories in a single request.

This XML file does not appear to have any style information associated with it. The document tree is shown below.

```
<?xml version="1.0"?>
<DESCRIPTION>
  Digitized First Byurakan Survey (DFBS) arvo.sci.am send questions to: aram_knyazyan@ipia.sci.am
</DESCRIPTION>
<DEFINITIONS>
  <COORDSYS ID="myj2000" system="eq_FK5" equinox="2000." epoch="2000."/ >
</DEFINITIONS>
<RESOURCE type="results">
  <INFO name="QUERY_STATUS" value="OK"/ >
  <PARAM name="pos" value="322.3,3" datatype="char" arraysizes="" unit="deg">
    <DESCRIPTION>Search position in form ra,dec in decimal degrees</DESCRIPTION>
  </PARAM>
  <PARAM name="size" value="1" datatype="double" unit="deg">
    <DESCRIPTION>Search diameter in decimal degrees</DESCRIPTION>
  </PARAM>
  <PARAM ID="param_max_records" name="max_records" datatype="int" value="50"/ >
  <PARAM ID="param_ra" name="ra" datatype="float" value="320.85874668 .. 322.85874668"/ >
  <PARAM ID="param_dec" name="dec" datatype="float" value="2 .. 4"/ >
  <PARAM ID="param_verb" name="verb" datatype="int" value="1"/ >
  <PARAM ID="param_mission" name="mission" datatype="char" value="dfbs_siap" arraysizes="" / >
  <PARAM ID="param_found_rows" name="found_rows" datatype="int" value="2"/ >
  <TABLE name="DFBS_SIAP">
    <DESCRIPTION>ArVO DFBS_SIAP Search: 2 row(s) returned!</DESCRIPTION>
    <FIELD ID="resourceid" name="resourceid" datatype="char" ucd="meta.id" arraysizes="">
      <DESCRIPTION>name of image file</DESCRIPTION>
    </FIELD>
    <FIELD ID="ra_j2000" name="ra_j2000" datatype="double" ucd="POS_EQ_RA_MAIN" unit="deg" ref="myj2000">
      <DESCRIPTION>RA from center of image array</DESCRIPTION>
    </FIELD>
    <FIELD ID="dec_j2000" name="dec_j2000" datatype="double" ucd="POS_EQ_DEC_MAIN" unit="deg" ref="myj2000">
      <DESCRIPTION>DEC from center of image array</DESCRIPTION>
    </FIELD>
    <FIELD ID="resourceurl" name="resourceurl" datatype="char" ucd="VOX:Image_AccessReference" arraysizes="">
      <DESCRIPTION>URL of image file</DESCRIPTION>
    </FIELD>
    <FIELD ID="id" name="id" datatype="char" ucd="ID_MAIN" arraysizes="">
      <DESCRIPTION>OPO-designated ID</DESCRIPTION>
    </FIELD>
    <FIELD ID="title" name="title" datatype="char" ucd="VOX:Image_Title" arraysizes="">
      <DESCRIPTION>General descriptive title</DESCRIPTION>
    </FIELD>
    <FIELD ID="subjects" name="subjects" datatype="char" ucd="meta.id" arraysizes="">
      <DESCRIPTION>Name of object(s) (semi-colon separated list)</DESCRIPTION>
    </FIELD>
    <FIELD ID="descriptions" name="descriptions" datatype="char" ucd="meta.node" arraysizes="">
      <DESCRIPTION>Full caption and related text.</DESCRIPTION>
    </FIELD>
    <FIELD ID="datasetid" name="datasetid" datatype="char" ucd="meta.dataset" arraysizes="">
      <DESCRIPTION>
        Sample list of datasets used to create image, with at least one dataset listed for each bandpass. Note this is not always a complete list of exposures used.
      </DESCRIPTION>
    </FIELD>
    <FIELD ID="referenceurl" name="referenceurl" datatype="char" ucd="meta.ref.url" arraysizes="">
      <DESCRIPTION>URL of press release images page at DFBSsite</DESCRIPTION>
    </FIELD>
    <FIELD ID="filesize" name="filesize" datatype="int" ucd="VOX:Image_FileSize" unit="byte">
      <DESCRIPTION>File size in megabytes</DESCRIPTION>
    </FIELD>
  </TABLE>
</RESOURCE>
```

Fig. 2. A result of request to ArVO SIA API.

ArVO SIA service is registered in European Virtual Observatory (EURO-VO) (Genova et al., 2015) registry and soon will be available in other astronomical registries.

ArVO is based on Armenian National Grid Infrastructure (ArmNGI) (Astsatryan et al., 2015) and uses its computational resources for making a simple images search. This allows making the response time of SIA API very quick.

4. Scientific applications of ArVO SIA service

So far, the First Byurakan Survey (FBS or Markarian Survey) has been completely digitized and DFBS database has been created for further scientific usage. Other ArVO digitization projects (Second Byurakan Survey—SBS, FBS blue stellar objects spectroscopy, variability of blazars, asteroids, etc.) are active including the digitization of the whole BAO Plate Archive with the creation of a full electronic database and interactive sky map (Mickaelian, 2016; Mickaelian et al., 2016a). All of this observing material and corresponding software form the basis of ArVO.

Opposite to almost all other astronomical observing material existing in the world, which are either in image or in spectral form, DFBS introduces a new type (as it was obtained with an objective prism and spectra for objects in the whole field are obtained): 2D images of the fields covered by the survey that include 2D spectra and 1D images of the extracted spectra for each object. This means that both SIA and SSA (implementation is planned) services are applicable and necessary for the DFBS. Besides the work with individual extracted spectra of objects (for further reduction, analysis, classification and study) very often there is a need to work with some fields around definite position, applying cone search.

Fig. 3 shows how efficient are DFBS 2D fields for study of objects having spectra for each of them compared to direct images where only the position, brightness and sometimes the morphology are available. 2D spectra give understanding on the nature of objects; they provide a preliminary classification and identification of non-optical sources, such as X-ray, IR and radio. That is why SIA service may be rather useful to have quick access to all necessary fields for further study.

DFBS and other 2D spectroscopic images may be used for the following tasks:

- Search for objects of definite types in the DFBS and other fields (extremely blue or extremely red objects, emission or absorption line objects, objects having flat or any other peculiar spectral energy distribution, SED);
- Optical identifications of X-ray, IR and radio sources, when the selection of the correct optical counterpart becomes reliable due to the understanding on the nature of the objects;
- Study of known asteroids and other moving objects and search for new ones.

SIA service is being widely used for access to many databases, however ArVO SIA service will be especially useful due to its application to the DFBS and other spectroscopic data.

5. Conclusion

The usage of the IVOA SIA protocol and computational resources of ArmNGI allow ArVO to provide a powerful astronomical data search API. The query parameters, which can be handled by SIA API, are updated and more search functionalities are regularly included




Fig. 3. A comparison of the same field from the Digitized Sky Survey (DSS; direct images) and DFBS (spectral images). 2D spectra significantly facilitate the search and identification of objects.

in it. SIA2.0 version is already a recommended standard and as ArVO SIA service is implemented on SIA1.0, upgrade to SIA2.0 is planned in the near future.

References

- Astsatryan, H., Sahakyan, V., Shoukourian, Y., Cros, P.-H., Dayde, M., Dongarra, J., Oster, P., 2015. Strengthening compute and data intensive capacities of Armenia. In: IEEE Proceedings of 14th RoEduNet International Conference Networking in Education and Research. pp. 28–33.
- Brunner, Robert, J., Djorgovski, S., George, Szalay, Alex, S., 2001. Virtual observatories of the future. In: *Astronomical Society of the Pacific Conference Series*, vol. 225.
- Dowler, P., 2016. Table Access Protocol Version 1.1, IVOA Recommendation.
- Genova, F., Allen, M., Arviset, C., Lawrence, A., Pasian, F., Solano, E., Wambsganss, J., 2015. Euro-VO Coordination of virtual observatory activities in Europe. *Astron. Comput.* 11, 181–189.
- Gudivada, V.N., Baeza-Yates, R.A., Raghavan, V.V., 2015. Big data: Promises and problems. *IEEE Comput.* 48 (3), 20–23.
- Mickaelian, A., 2016. Astronomical surveys and big data. *Balt. Astron.* 25, 75–88.
- Mickaelian, A., Nesci, R., Rossi, C., Weedman, D., Cirimele, G., Sargsyan, L., Gaudenzi, S., 2007. The digitized first Byurakan survey DFBS. *Astronom. Astrophys.* 464 (3), 1177–1180.
- Mickaelian, A.M., Andreasyan, H. V.A.H.R., Azatyan, N.M., Farmanyan, S.V., Gigoyan, K.S., Gyulzadyan, M.V., Khachatryan, K.G., Knyazyan, A.V., Kostandyan, G.R., Mikayelyan, G.A., Nikoghosyan, E.H., Paronyan, G.M., Vardanyan, A.V., 2016. BAO plate archive project: Digitization, electronic database and research programmes. In: *Astronomical Society of the Pacific Conference Series*, vol. 505. pp. 262–273.
- Mickaelian, A.M., Astsatryan, H.V., Knyazyan, A.V., Magakian, T.Y., Mikayelyan, G.A., Erastova, L.K., Hovhannisyan, L.R., Sargsyan, L.A., Sinamyan, P.K., 2016. Ten years of the armenian virtual observatory, astronomical surveys and big data. *Astron. Soc. Pac. Conf. Ser.* 505, 16–23.
- Murdin, P., 2000. Byurakan astrophysical observatory. *Encyclopedia Astronom. Astrophys.* 4.
- Dowler, P.F.B., Tody, D., 2015. Simple Image Access Specification Version 2.0, IVOA Recommendation.
- Plante, R., Williams, R., Hanisch, R., Szalay, A., 2008. Simple Cone Search Version 1.03, IVOA Recommendation 1.
- Quinn, P., Barnes, D., Csabai, I., Cui, C., Genova, F., Hanisch, B., Kembhavi, A., Kim, S., Lawrence, A., Malkov, O., Ohishi, M., Pasian, F., Schade, D., Voges, W., 2004. The international virtual observatory alliance: recent technical developments and the road ahead. In: *Proceedings of SPIE, Optimizing Scientific Return for Astronomy Through Information Technologies*, vol. 5493. 137–145.
- Tody, D., Dolensky, M., 2004. The simple spectral access protocol. In: *SPIE Astronomical Telescopes + Instrumentation*. 262–268.
- Tody, D., Plante, R., Harrison, P., 2009. Simple Image Access Specification Version 1.0, IVOA Recommendation.